

A note on scattering of electrons by atomic nucleus

H Ray

Department of Physics, S C College, Dhaniakhali, Hooghly-712 302, West Bengal, India

Received 16 September 1993, accepted 1 March 1994

Abstract : The proper radial wave function has been derived for the scattering of electrons from a bare nucleus. The asymptotic form of the total radial wave function required for evaluating scattering cross section has been computed with rigorous mathematical analysis. Point-like behaviour of scattering from nucleus at low and high energies has been pointed out.

Keyword : Elastic electron scattering, radial wave function, asymptotic form.

PACS Nos. : 25.30.Bf, 34.80.Bm

The scattering of electrons involving atomic nuclei and charged particles are of great interest in physics.

In the present problem, for example, the scattering of electrons by a bare nucleus has been considered. Let us assume that the energy of the incident electrons is sufficient to penetrate the electron-cloud surrounding the nucleus. An electron moving towards a nucleus experiences mostly the Coulomb-field around the nuclear region. The more energetic electron may either be scattered from the vicinity of the nuclear region or it may pass on through the nucleons and then get scattered.

The interaction potential outside the charged nuclear sphere is purely Coulombic and inside the nuclear sphere Coulomb field is modified. The effect of modification may be taken into account by a certain parameter α .

The wave function may be taken as [1]

$$U = \frac{1}{r} \sum_{l=0}^{\infty} c_l g_l(r) P_l(\cos\theta), \quad (1)$$

Acknowledgments

My sincere thanks are due to Prof N C Sil, Indian Association for the Cultivation of Science for going through the manuscript and to Prof T Roy, Jadavpur University for his guidance and checking up this work.

References

- [1] S Flugge *Practical Quantum Mechanics* (New York : Springer Verlag) I p 293 (1971)
- [2] S Flugge *Practical Quantum Mechanics* (New York : Springer Verlag) II p 274 (1971)